

Scale-dependent growth from a transition in dark energy dynamics

MUSTAFA A. AMIN¹, PHILLIP ZUKIN² AND EDMUND BERTSCHINGER³

*Dept. of Physics and Kavli Institute for Astrophysics and Space Research
Massachusetts Institute of Technology, Cambridge, MA 02139*

Abstract

We investigate the observational consequences of the quintessence field rolling to and oscillating near a minimum in its potential, *if* it happens close to the present epoch ($z \lesssim 0.2$). We show that in a class of models, the oscillations lead to a rapid growth of the field fluctuations and the gravitational potential on subhorizon scales. The growth in the gravitational potential occurs on timescales $\ll H^{-1}$. This effect is present even when the quintessence parameters are chosen to reproduce an expansion history consistent with observations. For linearized fluctuations, we find that although the gravitational potential power spectrum is enhanced in a scale-dependent manner, the shape of the dark matter/galaxy power spectrum is not significantly affected. We find that the best constraints on such a transition in the quintessence field is provided via the integrated Sachs-Wolfe (ISW) effect in the CMB temperature power spectrum. Going beyond the linearized regime, the quintessence field can fragment into large, localized, long lived excitations (oscillons) with sizes comparable to galaxy clusters; this fragmentation could provide additional observational constraints.

Two quoted “signatures” of modified gravity are a scale-dependent growth of the gravitational potential and a difference between the matter power spectrum inferred from measurements of lensing and galaxy clustering. Here, both effects are achieved by a minimally coupled scalar field in general relativity with a canonical kinetic term.

¹email: mamin@mit.edu

²email: zukin@mit.edu

³email: edbert@mit.edu

1 Introduction

Scalar fields are used ubiquitously in cosmology to provide a mechanism for accelerated expansion during both the inflationary [1, 2, 3] and current dark energy dominated epochs (for example, [4, 5, 6]). In the inflationary case, accelerated expansion occurs when the inflaton field is rolling slowly and it ends once the field starts oscillating around the minimum of its potential. The oscillatory phase is well studied and gives rise to rapid growth of inhomogeneities because of couplings to other fields or self interactions (see, e.g. [7, 8, 9, 10]). For a recent review, see [11]. Like the inflaton, the quintessence field provides a mechanism for accelerated expansion in the slow-roll regime. However, unlike inflation, the possibility of the accelerated expansion ending through quintessence oscillations is rarely discussed in the literature (however, see e.g. [12]). If quintessence was rolling slowly in the past, it is “natural” (though certainly not necessary) for it to enter an oscillatory regime as it finds its way towards a local minimum in its potential.

In this paper, we wish to understand the observational signatures of a transition to an oscillatory regime in the quintessence field, if such a transition occurs in our recent past ($z \lesssim 0.2$). An obvious observational signature is a late time change in expansion history. More importantly, as is the case with the inflaton, the coherent oscillations of quintessence near an anharmonic minimum are unstable and quickly fragment in a spatially inhomogeneous manner, giving rise to additional structure. This instability driven by self interactions of the field (the anharmonic terms) leads to a growth in structure that is much faster than the usual gravitational growth. For example, all potentials that have a quadratic minimum and are shallower than quadratic away from the minimum (see figure 1) suffer from this instability (in particular for wavenumbers $k \ll m$, where $m^2 \approx U''(\varphi)|_{\varphi \rightarrow 0}$. See, for e.g [13, 14, 9, 15]). We find that while the gravitational potential is influenced strongly by such growth in the quintessence fluctuations, the over density in baryons and dark matter does not change significantly. The scale dependent growth in the gravitational potential and the *absence* of similar growth in the matter power spectrum provides a signature to confirm or rule out such a transition in the dynamics of the quintessence field. In addition, if the quintessence field becomes nonlinear, it can fragment rapidly into long lived, localized excitations called oscillons (see e.g. [16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 9, 29, 30, 31, 32, 15]) with size $l \sim \text{few} \times m^{-1} \sim \text{few} \times \text{Mpc}^*$. Although we do not pursue this in detail in this paper, in appendix B, we comment on how such rapid nonlinear fragmentation of the field, and to a lesser extent, the time-dependent evolution of oscillon configurations themselves could have interesting ISW signatures.

While similar resonant phenomenon are commonly invoked in the early universe, for example during preheating ($z \gg 10^{20}$) [11], few observational signatures exist (e.g. [33, 34, 35, 36]). The reason for this lack of observational signatures on astrophysical scales is that resonance is effective only on subhorizon scales. Since high redshifts imply a small Hubble

*We will frequently relate the mass of the scalar field m , to length and time scales, without explicitly writing \hbar and c . We set $\hbar = c = 1$ throughout the paper.

horizon, only small astrophysically inaccessible scales today (of order meters for GUT scale inflation) become excited. Moreover, on these small scales, the thermal radiation dominated state of the universe at the time of big bang nucleosynthesis smooths out inhomogeneities. Since in our scenario the transition in quintessence is happening today, the resonance now happens on astrophysically accessible scales. As a result these models can be strongly constrained.

The observationally interesting phenomenology of quintessence oscillations comes at a price. In addition to the usual problems with quintessence models (such as the smallness of the energy density, long range forces, etc.), for the quintessence potentials we work with, we require that the transition to an oscillatory regime happens close to today. This imposes an additional tuning for the initial conditions of the field.

Related to the present paper [14], Johnson and Kamionkowski discuss the dynamical instability arising from a “small” anharmonic term in oscillating dark energy models and conclude that such instabilities render oscillating dark energy models unsuitable for providing a sustained cosmic acceleration period. While we rely on these very instabilities to source the rapid growth of the field fluctuations and the gravitational potential, we envision the transition happening late enough so that this instability does not significantly affect the observable expansion history. We then calculate and discuss in detail the additional observables that can be used to probe the rapid growth in the gravitational potentials arising from the dynamical instability. We reiterate, that we are not interested in the oscillatory phase providing the accelerated expansion.

We briefly mention a few papers which explore clustering in quintessence like fields, though through very different mechanisms. Observational effects of quintessence clustering caused by phenomenologically varying its speed of sound (*not* from resonant behavior discussed here) has been investigated by several authors (see, for example, [37, 38, 39, 40, 41, 42, 43]). Instabilities in coupled dark energy-dark matter models have also been considered in [44]. In addition, rapid transitions in dark energy, in the context of coupled/unified dark matter and dark energy models where dark energy “switches on” at late times have been explored by [45] (also see references therein). In our case, dark energy “turns off” at late times as the quintessence field starts oscillating, leading to a rapid growth in field fluctuations. With a somewhat different motivation, in [46], the authors discuss how rapid, extremely low redshift ($z \lesssim 0.02$) transitions in dark energy can be hidden from expansion history measurements. However, they did not consider the effects of such transitions on perturbations, which is the focus of this work.

In this exploratory paper, we restrict ourselves to single field quintessence models, assume a minimal coupling to gravity and impose no non-gravitational couplings to other fields. We include WIMP (Weakly Interacting Massive Particle) dark matter along with the quintessence field and assume a spatially flat universe. We limit ourselves to a linear treatment of the fluctuations, including scalar gravitational perturbations, except when we discuss the nonlinear fragmentation of the quintessence field and the formation of robust,

localized quintessence excitations (oscillons). We focus on a case that is consistent with expansion history and galaxy clustering but could potentially be ruled out by measurements of the gravitational potential, in particular via the CMB temperature power spectrum at large angular scales. Our intention is to stress the interesting phenomenology of such models and to point out that such transitions can be better constrained by going beyond the measurements of expansion history and growth of structure in galaxies alone.

The rest of the paper is organized as follows. In section (2) we introduce and motivate the form of the quintessence potential used in this paper. In section (3) we work through the evolution of the quintessence field in an FRW universe. We highlight initial conditions and important regimes of evolution as well as constraints placed on the parameters from observations of the expansion history. In section (4) we discuss the evolution of linearized fluctuations in the quintessence field, dark matter and the gravitational potential, with a special emphasis on their evolution during the oscillatory phase of the background quintessence field. In the same section we also discuss the domain of validity of our linearized treatment and comment on the nonlinear evolution of the field. In section (5) we compute observables such as lensing power spectra and the ISW contribution to the CMB temperature anisotropy. We present our conclusions in section (6). We also include two appendices. In appendix A, we provide some details of Floquet analysis and an algorithm used in this paper for calculating Floquet exponents. In appendix B, we provide an estimate of the ISW effect resulting from evolution of quasi/nonlinear quintessence fluctuations and oscillons.

2 The model

We consider a quintessence field governed by a potential of the form (see figure 1):

$$U(\varphi) = \frac{m^2 M^2}{2} \left[\frac{(\varphi/M)^2}{1 + (\varphi/M)^{2(1-\alpha)}} \right], \quad (1)$$

where $0 < \alpha < 1$.[†] This choice was motivated by monodromy and supergravity models of inflation [47, 48, 49, 50, 51] and a recent model of axion quintessence [52] ($\alpha = 1/2$). The potential has a quadratic minimum and for very large field values it asymptotes to a shallower than quadratic form:

$$\begin{aligned} U(\varphi) &\approx \frac{m^2}{2} \varphi^2 & \varphi \ll M, \\ U(\varphi) &\approx \frac{m^2 M^2}{2} (\varphi/M)^{2\alpha} & \varphi \gg M. \end{aligned} \quad (2)$$

The scale M determines where the potential changes shape whereas the scale m determines the curvature $U''(0)$ at the bottom of the well. We will consider cases where the field rolls

[†]We write φ instead of $|\varphi|$ to avoid clutter.




Figure 1: Initially, the field rolls slowly causing accelerated expansion of space. As it enters the oscillatory regime, the acceleration stops and a rapid, exponential growth of field fluctuations begins.

slowly for $\varphi \gg M$, behaving like dark energy and then enters an oscillatory regime with $\varphi \sim M$ after $z \sim 0.2$. When the field oscillates around the minimum, the “opening up” of the potential:

$$U(\varphi) - (1/2)m^2\varphi^2 < 0 \quad \varphi \neq 0, \quad (3)$$

leads to rapid, scale dependent growth of scalar field fluctuations via parametric resonance (see section (4)). In the nonlinear regimes, it leads to the formation of localized field excitations called oscillons [9, 29].

Similar phenomenology can arise in models such as pseudo-Nambu-Goldston-Boson (pNGB) quintessence (eg. [53])

$$U(\varphi) = m^2 M^2 [1 - \cos(\varphi/M)]. \quad (4)$$

For potentials (1) and (4), a limited range of parameters will reproduce the observed expansion history, allow for a few oscillations in the field close to today *and* allow for rapid growth of structure. For the pNGB like models, the slow roll dynamics of the field necessary for accelerated expansion, only occur if $M \sim m_{\text{Pl}}$ (unless $\varphi \rightarrow \pi M$). However, as we discuss in section (4), efficient resonance in an expanding universe requires $M \ll m_{\text{Pl}}$. As a result we do not get rapid growth of structure here, except in cases of extreme fine tuning of the initial conditions, and do not pursue this model any further in this paper.

For the potential in (1), we find the requirement of a few oscillations close to today translates into

$$10^2 H_0 \lesssim M, m \lesssim 10^{-2} m_{\text{Pl}} \quad \text{and} \quad \alpha \ll 1. \quad (5)$$

We study these constraints in more detail in the next section.

Finally, we note that resonant growth of fluctuations also arise when scalar fields oscillate in potentials with asymmetric minima

$$U(\varphi) \approx \frac{m^2}{2}\varphi^2 + \frac{M}{3}\varphi^3 + \dots,$$

and is possible in potentials with a non-quadratic minimum or in potentials which do not necessarily open up (though the band of resonant wave-numbers can be narrow). The key requirement is that there are anharmonic terms in the potential that provide a time varying, periodic frequency in the equation of motion for the fluctuations.

3 Homogeneous evolution

In this section, we describe how to choose initial conditions for the background quintessence field and parameters in the potential so that the field both oscillates at late times and reproduces the observed expansion history. We do this by trying to match the Λ CDM expansion history. This is more restrictive than directly using observations, but makes the following discussion more transparent.

The equations of motion for the homogeneous quintessence field are

$$\begin{aligned}\ddot{\varphi} + 3H\dot{\varphi} + U'(\varphi) &= 0, \\ H^2 &= \frac{1}{3m_{\text{Pl}}^2} \left[\frac{\dot{\varphi}^2}{2} + U(\varphi) \right] + H_0^2 \frac{\Omega_{\text{dm}}}{a^3},\end{aligned}\tag{6}$$

where Ω_{dm} is the fraction of the critical energy density in WIMPs today, H is the Hubble parameter and H_0 is its value today[‡]. The dot “.” stands for a derivative with respect to cosmic time and the scale factor $a = 1$ today. We ignore the contribution from radiation since we are only interested in late times.

Current observations are consistent with the quintessence field behaving like a cosmological constant in the recent past as well as during the matter dominated epoch [54]. This implies that the field’s equation of state satisfies $w \equiv (\dot{\varphi}^2 - 2U)/(\dot{\varphi}^2 + 2U) \sim -1$ (atleast for $z \gtrsim 0.2$), which occurs if the ‘slow roll’ condition $\dot{\varphi}^2 \ll U$ is satisfied. Imposing that the energy density in the quintessence field during matter domination has the same order of magnitude as the ρ_Λ in Λ CDM, we find:

$$\left(\frac{m}{H_0}\right)^2 \left(\frac{\varphi_i}{M}\right)^{2\alpha} \sim 6(1 - \Omega_{\text{dm}}) \left(\frac{m_{\text{Pl}}}{M}\right)^2\tag{7}$$

where φ_i is the initial value of the field deep in the matter dominated epoch. Note that for simplicity, we’ve assumed that $\varphi_i \gg M$ and later justify our assumption at the end of the analysis. Since we’re only interested in scaling relationships, we’ll ignore factors of order unity from now on. The above gives one constraint between parameters in the potential and the field’s initial conditions.

[‡]we choose the subscript “dm” for WIMPs instead of the more usual m to avoid confusion with the mass of the scalar field

Imposing oscillations in the field at late times gives a second constraint. To derive this, we solve equation (6) in the slow roll, matter dominated regime, assuming the field does not change significantly from its initial value. We find (assuming $\varphi_i \gg M$):[§]

$$\frac{\varphi}{M} - \frac{\varphi_i}{M} \sim \begin{cases} -a^3(m/H_0)^2(M/\varphi_i)^3 & \text{if } \alpha = 0, \\ -a^3(m/H_0)^2(M/\varphi_i)^{1-2\alpha}\alpha & \text{if } \alpha > 0. \end{cases} \quad (8)$$

Though the above equation of motion is not valid in the oscillation regime, it gives an approximate scale factor (a_*) when oscillations begin ($\varphi/M \sim 0$). We find:

$$a_* \sim \begin{cases} (H_0/m)^{2/3}(\varphi_i/M)^{\frac{4}{3}} & \text{if } \alpha = 0, \\ (H_0/m)^{2/3}(\varphi_i/M)^{\frac{2-2\alpha}{3}}\alpha^{-\frac{1}{3}} & \text{if } \alpha > 0. \end{cases} \quad (9)$$

The actual scale factor when oscillations begin (a_{osc}) differs from a_* but the two are monotonically related. Combining eqs. 7 and 9, we calculate explicit expressions for m and φ_i in terms of M and a_* . We have:

$$\begin{aligned} \frac{\varphi_i}{M} &\sim \begin{cases} a_*^{\frac{3}{4}}(m_{\text{Pl}}/M)^{\frac{1}{2}} & \text{if } \alpha = 0, \\ a_*^{\frac{3}{2}}(m_{\text{Pl}}/M)\alpha^{\frac{1}{2}} & \text{if } \alpha > 0. \end{cases} \\ \frac{m}{H_0} &\sim \begin{cases} (m_{\text{Pl}}/M) & \text{if } \alpha = 0, \\ a_*^{\frac{-3\alpha}{2}}(m_{\text{Pl}}/M)^{1-\alpha}\alpha^{-\frac{\alpha}{2}} & \text{if } \alpha > 0. \end{cases} \end{aligned} \quad (10)$$

As discussed in more detail below (section 4), M sets the strength of the resonance between the perturbed quintessence field and the background quintessence field. We therefore keep it as a free parameter. We choose a_* so that the actual numerically computed oscillations begin around $a_{\text{osc}} \approx 0.8$. Corrections to distance measurements that depend on cosmological parameters beyond H_0 are small for scale factors larger than $a_{\text{osc}} \approx 0.8$. Hence modifying the expansion history there will not interfere with measurements that probe the cosmological expansion. Last, we fix m and φ_i using equation (10). The value of $\dot{\varphi}_i$ follows from equation (8).

[§]The apparent discontinuity between the cases $\alpha = 0$ and $\alpha > 0$ disappears when the full expressions are used.




Figure 2: (a) Homogeneous field evolution: Note that the field is relatively constant until a_{osc} after which it starts oscillating with a decaying amplitude. (b) The Hubble parameter increases initially compared to its ΛCDM counterpart but once the oscillations in the field begin, it starts decreasing again. The small oscillations in the Hubble parameter reflect the oscillations in the quintessence field. (c) The equation of state parameter $w \approx -1$ until a_{osc} , reflecting the cosmological constant like behavior due to the slowly rolling field, but starts oscillating about $w = 0$ after a_{osc} , reflecting the approximately non-relativistic matter like behavior of the oscillating field. (d) The acceleration parameter q reveals a period of decelerated expansion in the matter dominated era, followed by a period of accelerated expansion as the slowly rolling quintessence field begins to dominate the energy density. This continues until a_{osc} , after which q starts oscillating due to the oscillations in the quintessence field.

In practice, when numerically evolving equation (6) according to the above choices of initial conditions and potential parameters, we find that in order to get agreement with the Λ CDM expansion history in the past as well as produce oscillations close to today, we require additional fine tuning of the initial conditions and parameters. The procedure described above results in a value of H_0 that is smaller than the Λ CDM value. This is because by assumption, the Hubble parameter for both Λ CDM and quintessence are identical. At late times, however, the oscillations in the quintessence field cause it to behave like dark matter, giving rise to a steeper fall off than the Λ CDM Hubble parameter. For better agreement with H_0 , we increase the value of m until H_0 agrees to better than a percent.

Performing the above iteration, we find that α affects whether or not the field starts oscillating at low redshift. This should be expected since the asymptotic slope of the potential determines the time it takes to transition from slow-roll to oscillations. We can calculate this approximate time from the derivative of the quintessence field when slow roll ends. Using the slow roll condition described above, the value of the quintessence field when slow roll ends is:

$$\varphi_e \sim \begin{cases} (m_{\text{Pl}}/M)^{1/3} M & \text{if } \alpha = 0, \\ \alpha m_{\text{Pl}} & \text{if } \alpha > 0. \end{cases} \quad (11)$$

Assuming $3H\dot{\varphi} \sim -U'$, taking this transition to happen close to today, and using the constraints in equation (10), we find the change in scale factor (Δa) between the beginning of oscillations ($\varphi \sim 0$) and the end of slow roll is given by:

$$\Delta a \sim \begin{cases} (M/m_{\text{Pl}})^{2/3} & \text{if } \alpha = 0, \\ \alpha^{1-\alpha} & \text{if } \alpha > 0. \end{cases} \quad (12)$$

The larger the α , the longer it takes for oscillations to start. Hence, consistency with a Λ CDM expansion history and having late time oscillations limits $\alpha \ll 1$ for the case of nonzero α and $M \ll m_{\text{Pl}}$ for vanishing α . For the rest of this paper, we specialize to $\alpha = 0$. In summary, we keep M and a_* as free parameters and choose m , φ_i and $\dot{\varphi}_i$ so that the value of H_0 agrees with the measured value, the field behaves like dark energy initially, and the field starts oscillating at $a_{\text{osc}} \approx 0.8$.

For the above prescription, we calculate the evolution of the quintessence field, the Hubble parameter normalized by Λ CDM's Hubble parameter, the field's equation of state parameter w and the acceleration parameter $q \equiv \ddot{a}a/\dot{a}^2$ as a function of scale factor. We show the results in figure 2 for

$$m_{\text{Pl}}/M = 500, m/H_0 = 1130.6, \varphi_i/M = 23.7 \text{ with } a_{\text{osc}} = 0.82. \quad (13)$$

Unless otherwise stated we will use these as fiducial parameter values throughout this paper and refer to them as fiducial parameters. As constructed, the field remains approximately

constant for $a \ll a_{\text{osc}}$, rolling slowly towards the minimum. Notice, as assumed initially, that $\varphi \gg M$ at early times. At $a_{\text{osc}} \sim 0.8$, the field starts to oscillate with a slowly decaying amplitude:

$$\varphi \approx \varphi_{\text{osc}}(a) \sin(\omega t + \Delta). \quad (14)$$

Note that when $\varphi \ll M$, $\varphi_{\text{osc}}(a) \propto a^{-3/2}$. Because of the anharmonic terms in the potential, the frequency of oscillation (ω) depends on the amplitude of the field and is given by:

$$\omega \approx m \left\{ \frac{2}{\pi} \frac{1}{\sqrt{1 + (\varphi_{\text{osc}}/M)^2} \mathcal{E} [-(\varphi_{\text{osc}}/M)^2]} \right\} \leq m, \quad (15)$$

where \mathcal{E} is the complete elliptic integral. As $\varphi \rightarrow 0$, $\omega \rightarrow m$. Note that in deriving the above expression for the period, we've assumed $\omega \gg H_0$ so that energy is approximately conserved during an oscillation.

The Hubble parameter matches Λ CDM at early times since both models have the same amount of dark matter (see figure 2(b)). The Hubble parameter must then increase relative to Λ CDM in order for H_0 to agree, since the quintessence model's Hubble parameter falls off more steeply than Λ CDM at late times when the field starts to behave like non-relativistic matter. The deviation, for the fiducial parameters, is at most 5%. The equation of state parameter behaves as expected in the slow roll regime ($w = -1$) and oscillates between 1 and -1 after a_{osc} as energy swaps between its kinetic and potential parts (figure 2(c)). Lastly, the acceleration parameter reveals a period of decelerated expansion deep in the matter dominated era, followed by a period of accelerated expansion as the slowly rolling quintessence field begins to dominate the energy density. After a_{osc} , the acceleration parameter starts oscillating around $q = -1/2$ (figure 2(d)).

Observational constraints on the expansion history at late times arise primarily from distance measurements. Given our above prescription, we expect good agreement between distances calculated in our quintessence model and Λ CDM, which we show explicitly below. The comoving distance is given by:

$$\chi(a) = \int_a^1 \frac{da}{a^2 H(a)}. \quad (16)$$

In figure 3, we plot the percent difference in $\chi(a)$ between quintessence and Λ CDM. As expected, deviations are smaller close to today, where the Hubble parameter was tuned to be identical to the observed H_0 , and at early times, where dark energy can be ignored. The largest deviations are $\sim 4\%$. Since all observed distances, like the luminosity distance and the angular diameter distance, differ from the comoving distance by factors of the scale factor, figure 3 implies that current observations cannot distinguish between the background evolution of quintessence and the background evolution in Λ CDM [55, 56, 57, 58, 59]. We have




Figure 3: The difference in comoving distance from an observer today, between the quintessence and Λ CDM models. Deviations are smaller today, where both cosmologies have the same H_0 , and at early times, where both cosmologies have the same amount of dark matter and dark energy.

restricted our analysis to $a_{\text{osc}} \sim 0.8$ because choosing $a_{\text{osc}} \ll 0.8$ gives rise to unacceptably large deviations from the observed expansion history.

4 Perturbation evolution

In this section we investigate the dynamics of the fluctuations in the quintessence field ($\delta\varphi$), the gravitational potentials and the overdensity in WIMPs (δ_{dm}) assuming linearized equations of motion. We ignore radiation and neutrinos, since we are interested in late time dynamics. We work in the Newtonian gauge where the metric is

$$ds^2 = -(1 + 2\Phi)dt^2 + a^2(1 - 2\Psi)dx^2, \quad (17)$$

and Φ and Ψ are the two scalar gravitational potentials. Tracking the evolution of the gravitational potentials and the two matter components (WIMPs and quintessence) requires two second order differential equations. Normally, they are taken to be the energy-momentum conservation equations for the matter components. The gravitational potentials can then be obtained from the Einstein equations (constraints). However, since the most interesting dynamics happen in the gravitational potentials and the field fluctuations, we work with them as the degrees of freedom and then obtain the overdensity in WIMPs from the constraints. The equations of motion for the quintessence field fluctuations and the gravitational potential are (in Fourier space):

$$\begin{aligned} \ddot{\delta\varphi}_k + 3H\dot{\delta\varphi}_k + \left[\frac{k^2}{a^2} + U''(\varphi) \right] \delta\varphi_k &= -2U'(\varphi)\Psi_k + 4\dot{\varphi}\dot{\Psi}_k, \\ \ddot{\Psi}_k + 4H\dot{\Psi}_k + \frac{1}{m_{\text{Pl}}^2}U(\varphi)\Psi_k &= \frac{1}{2m_{\text{Pl}}^2} \left[\dot{\varphi}\dot{\delta\varphi}_k - U'(\varphi)\delta\varphi_k \right]. \end{aligned} \quad (18)$$

The second equation is the diagonal, space-space component of the Einstein equations, where the right hand side is the pressure perturbation provided by the scalar field. Since the WIMP dark matter is assumed to be pressureless, it does not contribute to this source term. The second gravitational potential Φ_k does not appear in the above equations because there is no anisotropic stress at linear order for single, minimally coupled scalar fields. This makes the two gravitational potentials equal

$$\Psi_k = \Phi_k,$$

via the off diagonal space-space part of the Einstein equations.[¶] After evolving the equations for $\delta\varphi_k$ and Ψ_k , δ_{dm} follows from the time-time components of the Einstein equations:

$$\delta_{\text{dm}} = -\frac{a^3}{3H_0^2\Omega_{\text{dm}}} \left[\left(6H^2 - \frac{\dot{\varphi}^2}{m_{\text{Pl}}^2} + 2\frac{k^2}{a^2} \right) \Psi_k + 6H\dot{\Psi}_k + \frac{\dot{\varphi}}{m_{\text{Pl}}^2} \delta\dot{\varphi}_k + \frac{1}{m_{\text{Pl}}^2} U'(\varphi) \delta\varphi_k \right]. \quad (19)$$

4.1 Initial conditions and evolution during matter domination

We are interested in the behavior of $\delta\varphi_k$, Ψ_k and δ_{dm} for $a \gtrsim 0.8$. To solve (18) we need to specify, $\delta\varphi_k, \delta\dot{\varphi}_k, \Psi_k, \dot{\Psi}_k$ on some initial time slice. This can be done self-consistently using the Λ CDM solutions for Ψ_k and δ_{dm} , if the initial timeslice is chosen deep into the matter dominated era. During matter domination, the quintessence field is a small fraction of the total energy density. As a result, the $\delta\varphi_k$ do not contribute significantly to Ψ_k and we can use Ψ_k from a Λ CDM cosmology at these early times. We take these initial conditions for Ψ_k and $\dot{\Psi}_k$ directly from the output of CMBfast [60] at $a_i \approx 10^{-2}$ since by this scalefactor the anisotropic stress from neutrinos is negligible and the contribution of dark energy is yet to become important. We are then left with specifying the initial conditions in $\delta\varphi_k$, which requires understanding its evolution in the matter dominated era.

The evolution of adiabatic modes on superhorizon: $k/aH \ll 1$ scales is given by (see [61])^{||}

$$\delta\varphi_k = \Psi_k \frac{\dot{\varphi}}{H} \left[\frac{(H/a) \int_{t_i}^t a(t') dt'}{-1 + (H/a) \int_{t_i}^t a(t') dt'} \right]. \quad (20)$$

The term in the square brackets is constant during matter domination

$$\delta\varphi_k = -\frac{2}{3} \Psi_k \frac{\dot{\varphi}}{H}. \quad (21)$$

On subhorizon scales ($k \gg aH$), during the matter dominated era, the gravitational potential is determined by the fluctuations in the WIMP overdensity. Since Ψ_k is constant

[¶]For non-minimally coupled scalar fields, $\Phi_k - \Psi_k \propto f'(\varphi)/f(\varphi) \delta\varphi_k$ where $8\pi G_N \rightarrow f(\varphi)^{-1}$. In minimally coupled fields, the anisotropic stress is generated by second order terms: $\Phi_k - \Psi_k \propto \delta\varphi_k^2$.

^{||}We will ignore isocurvature modes in this paper

during matter domination, the equations of motion (18) become

$$\begin{aligned}\ddot{\delta\varphi}_k + 3H\dot{\delta\varphi}_k + \frac{k^2}{a^2}\delta\varphi_k &= -2U'(\varphi)\Psi_k, \\ \dot{\Psi}_k &= 0,\end{aligned}\tag{22}$$

where we have assumed $U''(\varphi)/H^2 \ll (k/aH)^2$. Under the assumption that $U'(\varphi)$ is slowly varying, the complete solution is

$$\delta\varphi_k = \frac{c_k}{k_H^3} \left[\cos(2k_H + \theta_k) + 2k_H \sin(2k_H + \theta_k) \right] - 2\Psi_k \frac{U'(\varphi)}{H^2} \frac{1}{k_H^2} \left[1 - \frac{7}{k_H^2} + \frac{35}{2k_H^4} \right], \tag{23}$$

where $k_H \equiv k/aH$ and c_k, θ_k are constants of integration set by initial conditions at the beginning of the matter dominated era. For solutions that are deep inside the horizon, $k_H \gg 1$ and only one term grows with time:

$$\delta\varphi_k \approx -2\Psi_k \frac{U'(\varphi)}{H^2} \frac{1}{k_H^2}. \tag{24}$$

The initial (transient) oscillatory behavior due to the homogeneous part of the solution 23 as well as the growth $\propto a^2$ for $a_i \ll a \ll a_{\text{osc}}$ due to the particular solution 24 can be seen in figure 5(a). We have also verified that although from equations (21) and (24), we see that $\delta\varphi_k$ grows as a^3 on superhorizon scales and as a^2 on small, subhorizon scales, their amplitude during matter domination does not become large enough to change the behavior of the potential Ψ_k .

With the equations (21) and (24) at hand, we set the initial conditions for all k_H using a piecewise interpolation between the subhorizon and superhorizon solutions (dashed curve in figure 6). Although we have taken care to faithfully characterize the asymptotic behavior of the initial conditions of $\delta\varphi$, in practice we find that changing the initial conditions by orders of magnitude does not affect our results significantly. This is because at late times, the particular solution of $\delta\varphi_k$ (determined by Ψ_k) dominates, significantly reducing the dependence on initial conditions.

4.2 Evolution during the quintessence dominated era

In this section, we investigate the dynamics of fluctuations during quintessence domination and find that the evolution differs before and after a_{osc} . We analyze these two regimes separately.

4.2.1 Before resonance $a < a_{\text{osc}}$

When $0.5 < a < a_{\text{osc}}$, the homogeneous quintessence field rolls slowly with an energy density larger than, but comparable, to the WIMP density. The behavior of $\delta\varphi_k$ on super-horizon

scales is still determined by (20), but H decreases more slowly than during the matter dominated era.

In the subhorizon regime, the quintessence perturbations grow faster as we approach a_{osc} because the field starts rolling more rapidly (see right hand side of equation (18)). However, these fluctuations are still not strong enough to prevent the decay of the gravitational potential, which is caused by a transition to a dark energy dominated epoch. During this epoch, the evolution of the Ψ_k is determined by H and does not show any resonant behavior. The behavior of the $\delta\varphi_k$ and Ψ_k discussed here can be easily see in figure 5(a) and (b) for $0.5 \lesssim a \lesssim a_{\text{osc}}$. The thin black line represents the evolution of the same mode of Ψ_k in Λ CDM.

4.2.2 Resonance $a > a_{\text{osc}}$

We now come to the most interesting era ($a > a_{\text{osc}}$) with regards to the evolution of fluctuations. In this era, the homogeneous field begins to oscillate with an amplitude dependent frequency $\omega < m$ (see equation (15)). This leads to a rapid growth in the field fluctuations for certain characteristic ranges of wavenumbers. To understand this, we initially ignore expansion ($H = 0, a = 1$) and the gravitational perturbations ($\Psi_k = 0$) in equations (18).

Resonance in Minkowski space

The equation of motion for $\delta\varphi_k$ then becomes:

$$\delta\ddot{\varphi}_k + [k^2 + U''(\varphi)] \delta\varphi_k = 0. \quad (25)$$

Since the homogeneous field φ is periodic in time, $U''(\varphi)$ is periodic as well as long as $U''(\varphi) \neq \text{constant}$, which occurs for potentials with anharmonic terms (as is the case with our potential in (1)). This yields an oscillator with a periodically varying frequency, whose solutions can be analyzed via standard Floquet methods (for example, see [62]). For the interested reader, we review the main aspects of Floquet analysis in the appendix. Under certain conditions (see appendix A), Floquet's theorem guarantees that the general solution to equation (25) can be written as

$$\delta\varphi_k(t) = e^{\mu_k t} P_+(t) + e^{-\mu_k t} P_-(t), \quad (26)$$

where $\pm\mu_k$ are the Floquet exponents and $P_{\pm}(t)$ are periodic functions with the same period as $U''(\varphi)$.** The Floquet exponent μ_k depends on the amplitude of the “pump” field φ_{osc} , as well as the wavenumber k . There exists an unstable, exponentially growing solution if the real part of the Floquet exponent $\Re(\mu_k) \neq 0$. In figure 4(a), we show $|\Re(\mu_k)|$ as a function

**When the Floquet exponents are *zero*, there exist another class of solutions $\delta\varphi_k(t) \propto tP_1(t)$ and $\delta\varphi_k(t) \propto P_2(t)$ where $P_{1,2}(t)$ are periodic functions.




Figure 4: In figure 4(a), we plot the magnitude of the real part of the Floquet exponent $|\Re(\mu_k)|$ as a function of the physical wavenumber k_p and the amplitude of the background field φ_{osc} . Yellow corresponds to a larger value than orange, whereas dark red regions have $\Re(\mu_k) = 0$. As the universe expands, the wavenumber of a given mode as well as the amplitude of the background field redshift. As a result, a given mode “sees” different Floquet exponents as it traverses the Floquet chart along the thin white lines. In figure 4(b) we show the Floquet exponent “seen” by a mode with a co-moving wavenumber $k \sim 0.05m$ (along the left-most dashed white line in Figure 4(a)). Modes with $k \gtrsim 0.5m$ pass through many resonance bands in a single oscillation of the homogeneous field and undergo stochastic resonance (see text).

of the amplitude φ_{osc} and wavenumber. The color represents the magnitude of the real part of the Floquet exponent. Yellow corresponds to a larger $|\Re(\mu_k)|$ than orange, whereas red corresponds to $\Re(\mu_k) = 0$. Without expansion, neither φ_{osc} nor the momentum k redshift with time. As a result, the evolution of modes is determined by the Floquet exponent at single point in the $(k, \varphi_{\text{osc}})$ plane. This no longer holds true in an expanding universe.

Resonance in an expanding universe

The equation of motion for $\delta\varphi_k$ in an expanding universe (still ignoring Ψ_k) is

$$\ddot{\delta\varphi_k} + 3H\dot{\delta\varphi_k} + \left[\frac{k^2}{a^2} + U''(\varphi) \right] \delta\varphi_k = 0. \quad (27)$$

To understand parametric resonance in an expanding background, we make the following identifications:

$$\begin{aligned} ka^{-1} &\rightarrow k_p, \\ \varphi_{\text{osc}}(a) &\rightarrow \varphi_{\text{osc}}. \end{aligned} \quad (28)$$

Here k_p is the physical wavenumber and $\varphi_{\text{osc}}(a)$ is the decaying envelope of the oscillating field. This identification (28) defines a trajectory in the $k_p - \varphi_{\text{osc}}$ plane (white lines in figure 4(a)). The approximate amount of amplification undergone by a given mode is obtained by integrating $|\Re(\mu_k)|$ along the corresponding trajectory in the $k_p - \varphi_{\text{osc}}$ plane. The $|\Re[\mu_k(t)]|$ as seen along such a trajectory [left-most dashed, white line in figure 4(a)] is shown in figure 4(b). Then, schematically, the evolution envelope of the amplified modes is (ignoring the oscillatory piece $P_{\pm}(t)$):

$$\delta\varphi_k(t) \sim \frac{\delta\varphi_k(t_i)}{a^\gamma(t)} \exp \left[\int_{t_i}^t d\tau |\Re[\mu_k(\tau)]| \right] = \frac{\delta\varphi_k(t_i)}{a^\gamma} \exp \left[\int^a d \ln \bar{a} \frac{|\Re[\mu_k(\bar{a})]|}{H(\bar{a})} \right] \quad (29)$$

where in the second equality we use the scalefactor a as a time co-ordinate and $\gamma = 3/2$ when $\varphi_{\text{osc}} \ll M$. We assume that the frequency of oscillation $\omega \gg H$. This expression (29) is meant to give intuition about the resonant behavior in an expanding universe and should be used with care, especially when multiple bands are involved since the phase of the oscillations can play a role when different bands are traversed.

When a mode traverses only the first resonance band, the fluctuations get a boost every time the homogeneous field crosses zero. This can be thought of as a burst of particle production. When a large number of bands are traversed within a single oscillation of the homogeneous field, we enter the regime of stochastic resonance. The amplitude of the mode still changes dramatically at zero crossings of the homogeneous field, however we are no longer guaranteed growth at every such instant. Although over longer time scales the fluctuations grow, at some zero crossings they can also decrease. For our scenario stochastic resonance is seen for modes with $k \gtrsim 0.5m$. A more detailed discussion of these different regimes can be found in [63].

For our scenario, the modes that grow the fastest are the ones with $k \ll m$ (ones traversing the first band). To get a sense of what is required of the parameters for these modes to grow rapidly, let us now concentrate on the right most expression in (29). If the argument of the exponent is significantly larger than one, then we will have rapid growth in fluctuations. Using $a_{\text{osc}} \approx 0.8$, resonance takes place in the logarithmic interval $\Delta \ln a \sim 0.2$. Hence, one gets rapid growth in fluctuations when $|\Re(\mu_k)|/H \gtrsim 10$. During the oscillatory regime $H \approx H_0 \sim m(M/m_{\text{Pl}})$ while from figure 4, the real part of the Floquet exponent seen by a mode in the first band has value of $|\Re(\mu_k)| \sim 0.1m$. Thus we need $M/m_{\text{Pl}} \lesssim 10^{-2}$ for efficient resonance. However, as we saw in section (3), in the class of models described by (1), consistency with the observed expansion history and the requirement of a few oscillations in the homogeneous field close to today, automatically yields $M/m_{\text{Pl}} \lesssim (\Delta a)^{3/2} \approx .03$ (equation (12)). Hence in such models, resonance is almost inevitable.

Importantly, this is a non-gravitationally driven growth, driven by the homogeneous, oscillating field $\omega \gg H$. Hence this growth can happen on a time scale which is *significantly shorter* than H^{-1} .

Resonance in an expanding universe including local gravity

Let us now include the gravitational potentials in the equations of motion and consider resonant phenomenon in the complete coupled system given by equation (18). When the field driven resonance is efficient, the gravitational potential Ψ_k does not play a significant role in the field dynamics. The evolution of Ψ_k , on the other hand, is affected significantly by the resonant growth in $\delta\varphi_k$ since the quintessence field dominates the energy density. We stress that Ψ_k can only be ignored in the evolution of $\delta\varphi_k$ for modes undergoing strong resonance, when the self interactions dominate over the gravitational one. In particular, as we have seen in the previous sections $\delta\varphi_k$ just before resonance is determined by Ψ_k , and as a result ignoring Ψ_k before resonance is not justified. In addition, the gravitational potential includes contributions from the WIMP dark matter overdensity via the constraint equations. These two considerations make the analysis of equation (18) in the resonance regime non-trivial and we have to rely on numerical solutions. A more detailed *analytical* analysis of resonance in an expanding universe including the effects of the gravitational potential will be pursued elsewhere. Below we discuss the numerical solutions during resonance in a bit more detail.

Typical, rapidly growing solutions for $\delta\varphi_k$ and Ψ_k with $k \approx 0.05m$ are shown in figure 5(a) and (b) respectively. Note the rapid, nearly exponential growth of Ψ_k and $\delta\varphi_k$ after $a_{\text{osc}} \approx 0.8$ (vertical dashed line). The thin black lines show the evolution of the same modes in Λ CDM. The growth in the gravitational potential is somewhat delayed compared to the field. This is because the energy density in the field has to first become comparable to that of dark matter. Before this happens, the gravitational potential does not experience scale dependent growth (though it still responds to the changing expansion history).




Figure 5: (a) Evolution of $\delta\varphi_k$ with $k \sim 0.05m \approx 0.01\text{Mpc}^{-1}$. During matter domination, $\delta\varphi_k \propto a^2$ and grows relatively slowly. However, after a_{osc} , the mode grows exponentially fast due to parametric resonance. At a_{nl} the fluctuations become nonlinear and linear evolution is no longer applicable. (b) Evolution of the scalar gravitational perturbation ($k \sim 0.05m$). Before resonance, the evolution is similar to that expected in usual slow roll quintessence (or ΛCDM , thin black line) model. It is constant deep in the matter dominated era and starts decaying as quintessence takes over. Unlike slow roll quintessence, after a_{osc} , the potential grows rapidly until a_{nl} . This leads to a scale dependent signal in observations that are sensitive to the gravitational potential. (c) The evolution of the WIMP overdensity is not significantly affected by the resonant growth. The departure from ΛCDM in this case is almost entirely due to the slight deviations in the expansion history. The normalization is set to one for all the above modes at $a = a_i$.

So far we have ignored discussing the evolution of the WIMP overdensity since it is determined from the constraint equation (19). Naively one might expect to see scale dependent departures in the behavior of δ_{dm} . However, despite the rapid growth in the gravitational po-

tential after a_{osc} , δ_{dm} does not deviate significantly from its ΛCDM counterpart (see 5(c)) in the linear regime. Although difficult to see from the constraint equation (19), this behavior can be understood by considering the conservation equation for the WIMPs:

$$\ddot{\delta}_{\text{dm}} + 2H\dot{\delta}_{\text{dm}} = -\frac{k^2}{a^2}\Psi_k + 3\left(2H\dot{\Psi}_k + \ddot{\Psi}_k\right). \quad (30)$$

Heuristically, we see that on subhorizon scales, δ_{dm} is obtained from a double time integral of the potential Ψ_k . This delays the response of the δ_{dm} to the Ψ_k . The small departure of δ_{dm} from its ΛCDM counterpart can be accounted for by the difference in expansion history between the quintessence and the ΛCDM models. In particular, since the Hubble parameter in our quintessence model is always slightly larger than its ΛCDM counterpart (see section 3), the growth of δ_{dm} in the quintessence model is slightly suppressed.

From the above discussion we see that the linearized fluctuations in the field and the gravitational potential grow rapidly. Eventually, the field fluctuations will become nonlinear and we cannot trust the linearized treatment. Hence it becomes important to understand, at least qualitatively, when the field becomes nonlinear as well as what happens thereafter. Although we discuss the nonlinearity of the scalar field fluctuations, we do not include the usual nonlinearity in δ_{dm} at late times. This nonlinearity in δ_{dm} is of course well studied, but to include it would take us too far beyond the scope of this paper.

4.2.3 Nonlinearity and oscillon formation

We cannot ignore the nonlinearity of the fluctuations when $U'(\varphi) \sim U''(\varphi)\delta\varphi$. In the oscillatory regime this happens at the scale factor a_{nl} when

$$\Delta_{\delta\varphi}(k, a_{\text{nl}}) \sim \varphi_{\text{osc}}(a_{\text{nl}}). \quad (31)$$

where $\Delta_{\delta\varphi}^2(k, a) = k^3 P_{\delta\varphi}(k, a)/2\pi^2$ with $\langle\delta\varphi_k\delta\varphi_{k'}\rangle \equiv (2\pi)^3 P_{\delta\varphi}(k)\delta^3(k-k')$ is the power spectrum of the field fluctuations (see figure 6(a)). Heuristically, the left hand side characterizes the mean square fluctuations over a spatial region of size L , ie.

$$\langle\delta\varphi^2\rangle_L^{1/2} = [\Delta_{\delta\varphi}(k, a)]_{k\sim L^{-1}} \quad (32)$$

In figure 6(b) the orange curves show the evolution of the (absolute value of) oscillatory homogeneous field and its envelope. The black curves show the maximum value of $\Delta_{\delta\varphi}(k, a)$ as a function of a and its envelope. The location where the two curves intersect $a_{\text{nl}} \approx 0.95$, is taken as the point beyond which the linearized equations cannot be trusted. In figure 6(a) we show $\Delta_{\delta\varphi}(k, a_{\text{nl}})$ (solid orange curve) and $\Delta_{\delta\varphi}(k, a_i)$. As mentioned in subsection 4.1, the normalization of $\Delta_{\delta\varphi}(k, a_i)$ is set by $\Delta_{\Psi}(k, a_i)$, which we take from WMAP 7 [55] ($\Delta_{\mathcal{R}}^2 = (9/25)\Delta_{\Psi}^2 = 2.42 \times 10^{-9}$ at $k = 0.002 \text{ Mpc}^{-1}$ and $n_s = 0.966$). We also note that although $k \sim 0.05 m \approx 0.01 \text{ Mpc}^{-1}$ is where we see the maximum deviation, resonance significantly enhances fluctuations for larger k as well, albeit through stochastic resonance.




Figure 6: (a) The orange curve shows the rms fluctuations in quintessence $\langle \delta\varphi^2 \rangle_{L \sim k^{-1}}^{1/2} = \Delta_{\delta\varphi}(k, a) = \sqrt{k^3 P_{\delta\varphi}(k, a) / 2\pi^2}$ at $a_{\text{nl}} = 0.95$, whereas the dashed curve is $10^3 \times$ the initial conditions at $a_i = 0.01$. Note the large bump due to resonant growth seen in the $\delta\varphi$ field at $k \approx 0.05 m \approx 0.01 \text{ Mpc}^{-1}$. The higher k regions undergo stochastic resonance. (b) To determine when the fluctuations become nonlinear (a_{nl}) we compare the rms fluctuations to the background field. The orange curves denote the homogeneous field whereas the black curves denote the maximum value of $\Delta_{\delta\varphi}(k, a)$ at each a . Note the enhancement at each zero crossing. Once the field fluctuations become nonlinear, different k modes couple and the field fragments into localized, long lived energy density configurations (oscillons).

In order to calculate the effect of the rapid growth of the gravitational potential on observables, we need its evolution until today, ie. $a \approx 1$. However, as discussed above, the nonlinearities of the field do not allow us to compute it for $a > a_{\text{nl}} = 0.95$. To remedy this, for the linearized calculation, we take a conservative approach and “freeze” the value of the potential at a_{nl} , setting $\Psi_k(a) = \Psi_k(a_{\text{nl}})$ for $a_{\text{nl}} \leq a \leq 1$. More realistically, the potential will evolve further in a scale-dependent manner as the scalar field perturbations undergo nonlinear evolution. Fragmentation of the scalar field (discussed below) will enhance the perturbations, at least on smaller scales (we provide some simple estimates of the ISW effect due to this nonlinear evolution in appendix B). Along with the break-down of linearized equations, we also note that after a_{nl} , large gradients in the scale field fluctuations will source anisotropic stress, violating our assumption of $\Psi = \Phi$.

Although in this paper we do not pursue nonlinear evolution of the scalar field fluctuations, below we digress slightly and point out some of the interesting phenomenology that results. As nonlinearity sets in, the evolution of different k modes becomes coupled, and

back-reaction from the perturbations curtails the resonant growth of fluctuations. The homogeneous field then fragments rapidly. For the type of potentials considered here (those having a quadratic minimum and a shallower than quadratic form away from the minimum), most of the energy density can eventually end up in localized, oscillatory, long lived configurations of the field called oscillons (e.g. [16, 18, 29]). The central density of these oscillons can be greater than the background energy density and their sizes are of order a few $m^{-1} \approx 10^{-3} H_0^{-1} \approx 4 \text{ Mpc}$. Although oscillons radiate energy through scalar radiation (e.g. [17, 64, 65, 30]), our limited radial simulations of individual oscillons reveal a lifetime of $\sim 10^6 m^{-1} \sim 10^3 H_0^{-1}$ for an oscillon with width of a few m^{-1} and field amplitude of order M . This makes them effectively stable compared to current cosmological timescales. Unlike the usual oscillons (eg.[16, 18, 65, 29]) which have a relatively stationary energy density profile, the oscillons in our model have an energy density that breathes in and out (also see [15]) at about twice the field oscillation frequency.

This phenomenon of oscillon production has been studied in the context of the early universe, in particular at the end of a similar scalar field driven inflationary period (e.g. [32, 10, 9, 15]). Here, we point out that it will likely happen at the end of the current period of cosmic acceleration as well. As with the early universe studies, we expect the oscillons to dominate the energy density. However, unlike the early universe, these oscillons have sizes which make them astrophysically accessible, with novel signatures such as the existence of dark, low redshift clusters with time dependent configurations made of the quintessence field. We stress that it is not guaranteed that the universe will become oscillon dominated before today. Although for the fiducial parameters, the field fluctuations will become nonlinear before today, it can take some time for the nonlinear field to enter a state where it is dominated by oscillons. This time can be longer than the time between a_{nl} and today. Quantifying this process, requires simulating the full nonlinear dynamics of the quintessence field, including nonlinearities of the WIMP overdensity, which is beyond the scope of this paper. Given the similarity of the potential with [15], we expect many of the qualitative results to carry over. However, to perform a reliable calculation to be compared with observations, one cannot completely ignore the gravitational perturbations, which makes the simulations somewhat more challenging.

5 Power spectra and observables

With our understanding of the evolution of individual modes, and the validity of the linearized equations, we now compute the power spectra of the gravitational potential $\Delta_\Psi^2(k, a) \equiv k^3 P_\Psi(k, a)/2\pi^2$ and the WIMP overdensity $P_{\delta_{\text{dm}}}(k, a)$ and their impact on observables such as galaxy clustering, lensing and the CMB.

The calculated power spectra $\Delta_\Psi^2(k, a_{\text{nl}})$ and $P_{\delta_{\text{dm}}}(k, a_{\text{nl}})$ (with $a_{\text{nl}} = 0.95$) are shown in figures 7 (thick orange lines). The thin black lines show the corresponding power spectra for the ΛCDM cosmology. The gray dashed lines show the power spectra at $a_i = 10^{-2}$. Note




Figure 7: (a) Power spectrum of the gravitation potential (orange). The dashed line denotes the initial conditions at $a_i = 0.01$ whereas the thin black line shows the Λ CDM power spectrum today. Notice the scale dependent growth of the gravitational potential for $k \sim 0.05m = 0.01\text{Mpc}^{-1}$, aside from which the potential decays in a manner similar to its Λ CDM counterpart. (b) The WIMP power spectrum (orange) does not show any features of the scale dependent growth, since the WIMPs do not have enough time to respond to the change in the gravitational potential. The difference from the Λ CDM counterpart (black) is primarily due to the difference in expansion history between the quintessence and Λ CDM models. The dashed line represents $10^3 \times P_{\text{dm}}(k, a_i)$. Note that since we are using δ_{dm} in the Newtonian gauge, $P_{\delta_{\text{dm}}}(k, a_i)$ shows an “upward turn” at small k values.

that since we are using δ_{dm} in the Newtonian gauge, $P_{\delta_{\text{dm}}}(k, a_i)$ shows an “upward turn” at small k values. By construction, the initial conditions in the power spectra $\Delta_\Psi(k, a_i)$ and $P_{\text{dm}}(k, a_i)$ in our quintessence model and Λ CDM agree. The scale dependent departures in $\Delta_\Psi(k, a)$ arise after $a_{\text{osc}} \approx 0.8$. As discussed previously, the enhancement can be trusted until $a_{\text{nl}} = 0.95$, after which we freeze the potential until $a = 1$. The wavenumber where the maximum departures are seen is $k \sim 0.05m \approx 0.01\text{Mpc}^{-1}$. For $k \ll H_0$ and $k \gtrsim m$, the final spectrum closely resembles the Λ CDM case. For fixed a_{osc} , we have checked that the magnitude of departure from Λ CDM increases with increasing m . This is consistent with the notion larger m corresponds to a larger m_{Pl}/M (see equation (10)) leading to more efficient resonance. As discussed for single modes, the potential $\Delta_\Psi(k, a)$ show dramatic, scale dependent departures from Λ CDM but the WIMP overdensity power spectrum does not.

Assuming a model for the bias, the WIMP power spectrum can be probed by measuring the two point correlation function of galaxies. Given the lack of scale dependence, and the

small difference ($\lesssim 6\%$) with Λ CDM we expect the departures from Λ CDM to be difficult to detect. ^{††} The fluctuations in quintessence are not directly observable, except through their gravitational imprint. This leaves us with the gravitational potential as the key probe to look for departures from Λ CDM. The rapidly changing potential in the quintessence model and the large scale dependent departures at low redshifts could affect the low multipoles of the CMB temperature and weak lensing power spectrum. With this in mind, we now calculate the weak lensing (convergence) power spectra and the CMB temperature anisotropy. We remind the reader that the differences that arise here are inspite of having made the expansion history consistent with current observations. An expansion history and galaxy power spectrum consistent with Λ CDM but scale dependent deviations in the lensing and CMB power spectra provides a unique probe of a late time transition in the quintessence field.

5.1 Integrated Sachs-Wolfe (ISW) effect

When CMB photons travel through an evolving gravitational potential, their energy changes (the ISW effect [66]). Since the rapid growth in the quintessence fluctuations induces large changes in the potential, we expect imprints of the late time quintessence transition on the CMB power spectrum through the ISW effect.

The angular power spectrum of the CMB temperature can be written as [67]

$$C_l = 4\pi \int d \ln k D_l^2(k) \Delta_\Psi^2(k)|_{\text{prim}} \quad (33)$$

where $\Delta_\Psi^2(k)|_{\text{prim}}$ is the primordial power spectrum. In the above, $D_l(k) \equiv \Delta_l(k)/\Psi_k^{\text{prim}}$, where $\Delta_l(k)$ on large angular scales (under the assumptions of adiabatic initial conditions and instantaneous recombination) is given by:

$$\begin{aligned} \Delta_l(k) &= \Delta_l^{\text{SW}}(k) + \Delta_l^{\text{ISW}}(k) \\ &\approx \left[\frac{1}{3} j_l(k\chi_e) + \frac{2}{3} \frac{k}{a_e H_e} j'_l(k\chi_e) \right] \Psi_k(a_e) + \int_{a_e}^1 da j_l(k\chi_a) \partial_a [\Psi_k(a) + \Phi_k(a)]. \end{aligned} \quad (34)$$

In the above expression, the subscript “e” stands for emission (ie. surface of last scattering), $\chi_a = \int_a^1 da/a^2 H(a)$ is the co-moving distance and $j_l(x)$ are spherical Bessel functions. The second term is the ISW term, which gets large if the potential evolves significantly between last scattering and today. In Λ CDM the ISW term gets a small contribution just after recombination since we are not quite matter dominated at that time. But, the main contribution on the scales of interest comes from late times as the universe starts becoming Λ dominated and the potentials start decaying. The same effect is also present in the quintessence model. However, there is an additional contribution from the late time rapid growth of the potential. This growth occurs in spurts (see figure 5(b)) and we can approximate $\partial_a \Psi_k$ as a

^{††}Note that this is reasonable since, like the WIMPs, galaxies will not have enough time to respond to the late time scale dependent quintessence fluctuations.

Dirac-delta function at the location of the jumps in the gravitational potential. As a result, the contribution of each jump to the ISW term can be easily evaluated:

$$^{(j)}\Delta_l^{\text{ISW}}(k) \approx 2j_l(k\chi_{a_j})\Delta\Psi_k(a_j), \quad (35)$$

where a_j is the scale factor where the potential jumps. For a jump of order a few Ψ_k , the “jump” term is larger than the smooth ISW contribution in Λ CDM. Note that there can be multiple jumps, both positive and negative, depending on the k mode in question. Also note that apart from the magnitude of the jump, the location a_j also plays a role by fixing the argument of the Bessel function.

We expect that this growth affects the low multipole moments of the CMB since the changes in the potential occur very recently. More specifically, the multipole range where we expect deviations is $l \lesssim k\chi_{a_j}$ where $k \sim 0.05m \approx 0.01\text{Mpc}^{-1}$. For the fiducial parameters chosen here, this yields $l \lesssim \text{few}$. Note that along with k , it is the smallness of χ_{a_j} that limits the effect to large angular scales.

In order to quantitatively calculate the effect of the quintessence perturbations on the CMB power spectrum, we use the $\Psi_k(a)$ and $\Phi_k(a)$ evaluated from CMBFAST [60] for $a < a_i = 0.01$. This captures the contributions from early ISW as well as early anisotropic stress (which we ignore at late times). For $a > a_i$ we use $\Psi_k(a)$ computed from our own independent code for Λ CDM and the quintessence model. As discussed before, for the quintessence case we set $\partial_a\Psi_k = 0$ for $a > a_{\text{nl}}$. With this entire solution at hand (for $a_e < a < 1$) we compute the $\Delta_l(k)$ and C_l . The ratio of the angular CMB power spectra for the quintessence and Λ CDM models is shown in figure 8(a). Note that this is large enough to potentially rule out this choice of parameters! Thus, we see that inspite of the effect being on large angular scales, ISW provides an excellent probe for constraining the considered transition in quintessence. One can change parameters, for example a_{osc} or m (equivalently M) to get the ISW effect small enough so that it is consistent with observations. One has to first make sure that such changes are still consistent with the expansion history (see discussion in section 3). Increasing m or a_{osc} or varying m and a_{osc} in opposite directions can lead to an expansion history consistent with observations. Let us now look at perturbations. In general, for a fixed a_{osc} , as we increase m (equivalently decrease M), the fluctuations grow more rapidly, shifting the time when the potentials grow rapidly to smaller a values. One might expect that this will make the ISW contribution even larger. However, one finds that the effect on $\Delta_l(k)$ is not quite as simple. First the change in growth of potential is quite sensitive to the number of zero crossing of the homogeneous field. In addition, changing parameters changes the wavenumbers which grow the fastest and the time when there is a rapid growth in the potential. Heuristically, this affects the argument of the $k\chi$ of the non-monotonic spherical Bessel function in $\Delta_l(k)$. Similar considerations apply to changing a_{osc} . As a result, it is somewhat difficult to apriori predict which combination of parameters (consistent with the observed expansion history) will also yield an acceptable ISW term. Indeed we find that $m \approx 2 \times 10^3 H_0$ and $a_{\text{osc}} \approx 0.82$ is entirely consistent with observations of the expansion history as well as the CMB. We also find regions of parameter space with smaller values of m but larger values of a_{osc} compared to the fiducial model consistent with




Figure 8: (a) Ratio of CMB temperature power spectrum at low multipoles between our quintessence and Λ CDM models. The large difference is due to the rapidly evolving gravitational potential in the quintessence model after a_{osc} (b) Ratio of the lensing (convergence) power spectrum at low multipoles between our quintessence and Λ CDM models. The departures are restricted to low l multipoles due to the proximity of the transition in the quintessence field responsible for the rapid growth in the gravitational potential. For lensing, the model does not agree with Λ CDM at large l due to the different expansion histories.

observations. A full sweep of parameters (including α) to determine regions consistent with observations is beyond the scope of this paper, but is certainly worth pursuing.

Before we move on to the calculation of the lensing power spectrum, let us briefly comment on the additional ISW that can result from nonlinear field evolution after a_{nl} , ignored so far in the linearized treatment. As noted in section 4.2.3, the rapid growth of the linearized scalar field perturbations is curtailed once the field perturbations become nonlinear (around a_{nl}). However, there will be further nonlinear evolution and rapid fragmentation of the field. To get an estimate of the resulting ISW effect, consider a spherical perturbation with mass \mathcal{M} and radius R , collapsing at speed v . Such a perturbation yields a temperature decrement of order $\sim G\mathcal{M}v/R$. For a sphere with an initial radius $R_{\text{nl}} \sim k_{\text{nl}}^{-1}$ (with $k_{\text{nl}} \sim 0.05m$), an initial density comparable to the background energy density and $v \sim 0.05$ (the approximate group velocity of a perturbation with wavenumber k_{nl}), the decrement can be $\sim \text{few} \times 10^{-5}$ for our fiducial set of parameters ($m \sim 10^3 H_0$). The signal is qualitatively similar to the Sunyaev-Zel'dovich (SZ) temperature decrement from galaxy clusters [68]. However unlike \lesssim arcmin angular scale of the SZ decrement, here, the angular scale is ~ 30 degrees. This simple estimate shows that it might be possible to get additional constraints on the quintessence transition from the ISW effect due to the nonlinear field evolution. Individual oscillations,

modeled by density configurations with time dependent radii $\sim \text{few} \times m^{-1}$ can also yield an additional ISW signal on smaller scales. We discuss a toy model for estimating the ISW signal from such nonlinear perturbations in appendix B. However, we again caution the reader that these numbers should be checked with input from detailed simulations of the nonlinear field dynamics including gravity.

5.2 Weak lensing

Observational constraints from weak lensing observations are often presented in terms of the weak lensing convergence (angular) power spectrum [69, 70]:

$$C_l^\kappa = 8\pi^2 \int_0^{\chi_e} d\chi W^2(\chi) \frac{l}{\chi} \Delta_\Psi^2(k = l\chi^{-1}, \chi) \quad (36)$$

where:

$$W(\chi) = \int_\chi^{\chi_e} d\chi' \frac{\chi' - \chi}{\chi'} \eta(\chi'), \quad (37)$$

$\Delta_\Psi^2(k, \chi)$ is the power spectrum of the gravitational potential and $\eta(\chi)$ is the radial distribution of sources, normalized to $\int \eta(\chi) d\chi = 1$. We use the source distribution:

$$\eta(z) \propto z^2 \exp[-(1.41z/z_{\text{med}})^{1.5}] \quad (38)$$

with $z_{\text{med}} = 1.26$. This distribution approximates the galaxy redshift distribution of the COSMOS survey[71]. As with the ISW effect, because of the closeness of the transition to the present day, we expect the signal to be largest at low multipoles $l \lesssim k\chi(a_{\text{nl}}) \sim \text{few}$. The proximity of the transition also picks out sources at approximately $2\chi(a_{\text{nl}})$. As a result as $a_{\text{nl}} \rightarrow 1$ ($\chi \rightarrow 0$), we run out of sources to be lensed. Thus in general, for a fixed enhancement of the gravitational potential, we expect the largest lensing signal to arise from the smallest a_{nl} still consistent with the expansion history.

The ratio between the weak lensing convergence power spectrum for the quintessence model and Λ CDM is shown in figure 8(b). The errors associated with measuring the convergence power spectrum is given by [69]:

$$\Delta C_l^\kappa = \sqrt{\frac{2}{(2l+1)f_{\text{sky}}}} \left(C_l^\kappa + \frac{\langle \gamma_{\text{int}}^2 \rangle}{\bar{n}} \right) \quad (39)$$

where f_{sky} is the fraction of the sky covered by the survey, \bar{n} is the measured galaxy number density, and $\langle \gamma_{\text{int}}^2 \rangle^{1/2} \approx .4$ is the galaxy intrinsic rms shear in one component. Using

survey parameters characteristic of LSST ($f_{\text{sky}} \approx .48$, $\bar{n} \approx 5.9 \times 10^8 \text{sr}^{-1}$) [72], we find $\Delta C_l^\kappa / C_l^\kappa \approx \{.9, .26\}$ for $l = \{2, 30\}$. Hence this particular quintessence model cannot be constrained using LSST measurements of the weak lensing power spectrum. The weak lensing convergence power spectrum for the quintessence model does not asymptote to the ΛCDM prediction at higher l because there is a difference in expansion history between the two models. Even this deviation would be difficult to see using LSST since the minimum $\Delta C_l^\kappa / C_l^\kappa \approx 0.1$ at $l \sim 300$.

6 Conclusions

For a slowly rolling quintessence field, it is natural (though not necessary) that the field will eventually start oscillating as it approaches a minimum in its potential. In this paper we have analyzed the consequences on the expansion history and structure formation of a quintessence field, which initially behaves like dark energy and starts oscillating around the minimum in its potential at late times ($a \gtrsim 0.8$). The potentials considered in detail here have a quadratic minimum and are shallower than quadratic away from the minimum. When a spatially homogeneous scalar field (φ) oscillates about the minimum of an such an anharmonic potential, it can pump energy into its spatially inhomogeneous perturbations ($\delta\varphi$) through parametric resonance. The amount of energy transfer depends on both the wavenumber of $\delta\varphi$ as well as the number of oscillations that take place in the background field. This leads to a rapid fragmentation of the homogeneous field and rapid resonant growth of the field fluctuations on time scales significantly shorter than H_0^{-1} .

In order to avoid discrepancies with the measured expansion history, and to simultaneously produce oscillations in the field, we have given a prescription for setting the initial conditions of the quintessence field and parameters in the potential. We have also explicitly shown that the potentials must be close to constant during the phase where the field is slowly rolling. Potentials that don't satisfy this requirement have too much of a delay between the end of slow roll and the beginning of oscillations, thus avoiding the rapid resonant growth of structure.

Given a model that produces a background expansion history in good agreement with the measured expansion history, we have shown how the gravitational potential and the overdensity in WIMPs is affected by the resonant growth of the field fluctuations. We found that the metric perturbations develop scale-dependent growth, with the scale set by the mass of the scalar field potential $m \sim \sqrt{U''(\varphi \rightarrow 0)} \sim 10^3 H_0$ ($k \lesssim 0.1m$). On the other hand, the dark matter overdensity remains featureless and very similar to the ΛCDM solution aside from an overall slight suppression because of the small difference in expansion history between ΛCDM and the quintessence model. This is because, the dark matter doesn't have time to respond to the changing potential. Note that scale dependent changes in the gravitational potential are normally attributed to modified gravity (e.g. [73, 74, 75]). Here however, we have shown that such a change can occur in general relativity with a minimally coupled

quintessence field with a canonical kinetic term.

The rapid growth of the potential significantly affects the ISW contribution to the temperature angular power spectrum of the CMB. For a range of parameters this yields the strongest constraint on such quintessence transitions. Since the metric perturbation develops a scale dependent change, the weak lensing power spectrum also offers a possible way to constrain this quintessence scenario, where dark energy undergoes a late time transition described above. Unfortunately, because of the proximity of the decay in dark energy, deviations from Λ CDM occur at low l multipole moments. A full treatment, which includes nonlinearities in the quintessence field, however, could give rise to deviations in the weak lensing and temperature power spectrum, at larger l values.

The nonlinear dynamics of the quintessence field would give rise to a wealth of new phenomenon including field fragmentation and possible formation of localized scalar field lumps which could provide additional observational constraints. The full nonlinear analysis, combining N-body simulations for the dark matter and lattice simulations for the scalar field, is beyond the scope of this paper, but provides a promising avenue to explore quintessence transitions and their consequences further. Recall that in spite of its strong clustering properties, our quintessence field is minimally coupled with a canonical kinetic term and does not have non-gravitational couplings to WIMP dark matter. This could make such models easier to simulate than models where gravity is modified (e.g. [76, 77, 78]), the quintessence field has non-gravitational couplings to dark matter (e.g. [79]), non-minimally coupled quintessence (e.g. [80]) or when non-standard kinetic terms are present.

In summary, if dark energy changes its nature close enough to the present time, it is possible to miss it in the expansion history measurements. For the models considered here, we have demonstrated that such a transition can dramatically change the gravitational potential power spectrum in a scale dependent way but leave the galaxy clustering unaffected, thus providing a possibly unique signature of such a transition. The best constraints on such transitions likely come from the ISW effect, followed by lensing. We expect, a more involved analysis, will provide additional constraints on such a transition once we include: (i) the nonlinear collapse in WIMPs and in the quintessence field, which will increase power on smaller scales (ii) couplings to other fields (ignored here). It would also be interesting to explore similar resonant growth in models with multiple ultralight scalar fields (not necessarily dark energy) motivated in [81] and studied in more detail in the context of structure formation by [82].

7 Acknowledgements

We would like to thank Alan Guth, David Shirokoff, Raphael Flauger, Richard Easther, Andrew Liddle, Volker Springel, Robert Crittenden, Marco Bruni, Ignacy Sawicki, Jonathan Pritchard, Navin Sivanandam and Charles Shapiro for useful conversations. We would espe-

cially like to thank Mark Hertzberg for comments on an early draft of this work and Antony Lewis for repeated and prompt help when we were trying to check our ISW calculation using CAMB. MA is supported by a Pappalardo Fellowship at MIT.

8 Appendix A

Floquet's theorem and calculating Floquet exponents

The linearized equations of motion for the fluctuations, neglecting the Hubble expansion and metric perturbation, are (in Fourier space)

$$\partial_t^2 \delta\varphi_k + [k^2 + U''(\bar{\varphi})] \delta\varphi_k = 0, \quad (40)$$

where k is the wavenumber. Since the homogeneous field φ is oscillating, $U''(\varphi)$ is periodic in time. This results in a linear system with periodic coefficients, which can be analyzed with Floquet theory. Floquet's theorem is most elegantly written in matrix form. Converting our second order equation of motion into a first order matrix equation, we find:

$$\partial_t x(t) = E(t)x(t), \quad (41)$$

where

$$x(t) = \begin{pmatrix} \delta\varphi_k \\ \partial_t \delta\varphi_k \end{pmatrix},$$

and

$$E(t) = \begin{pmatrix} 0 & 1 \\ -k^2 - U''(\bar{\varphi}) & 0 \end{pmatrix}.$$

Before stating Floquet's theorem, we need one more definition. The fundamental matrix solution, $\mathcal{O}(t, t_0)$, of equation (41) satisfies:

$$\begin{aligned} \partial_t \mathcal{O}(t, t_0) &= E(t)\mathcal{O}(t, t_0), \\ \mathcal{O}(t_0, t_0) &= 1. \end{aligned} \quad (42)$$

The fundamental matrix solution evolves the initial conditions $x(t_0)$ in time:

$$x(t) = \mathcal{O}(t, t_0)x(t_0). \quad (43)$$

Explicitly, $\mathcal{O}(t, t_0)$ consists of 2 columns that represent 2 independent solutions x_1, x_2 , which satisfy $x_1(t_0) = (1, 0)$ and $x_2(t_0) = (0, 1)$. Since $\det \mathcal{O}(t, t_0)$ is the Wronskian of linearly independent solutions and $\det \mathcal{O}(t_0, t_0) = 1$, we have $\det \mathcal{O}(t, t_0) = 1$.

We are now ready to state **Floquet's theorem** (without proof):

Consider the linear system

$$\partial_t x(t) = E(t)x(t), \quad (44)$$

where x is a column vector and E is a real, 2×2 matrix satisfying $E(t+T) = E(t)$ for all t . The fundamental solution $\mathcal{O}(t, t_0)$ is

$$\mathcal{O}(t, t_0) = P(t, t_0) \exp[(t - t_0)\mathcal{M}(t_0)], \quad (45)$$

where $P(t+T, t_0) = P(t, t_0)$ and $\mathcal{M}(t_0)$ satisfies $\mathcal{O}(t_0+T, t_0) = \exp[T\mathcal{M}(t_0)]$. The eigenvalues $\mu_{1,2}$ of $\mathcal{M}(t_0)$ are called Floquet exponents.

Since $\det \mathcal{O}(t_0 + T, t_0) = 1$, we have $\mu_1 + \mu_2 = 0$. Suppose, from now on, that $\mu_1 = -\mu_2 = \mu$. If $\mathcal{M}(t_0)$ has two linearly independent eigenvectors $e_{\pm}(t_0)$ corresponding to $\pm\mu$ (including $\mu = 0$), then the general solution can be written as

$$x(t) = c_+ \mathcal{P}_+(t, t_0) e^{\mu(t-t_0)} + c_- \mathcal{P}_-(t, t_0) e^{-\mu(t-t_0)} \quad (46)$$

where $\mathcal{P}_{\pm}(t, t_0) = P(t, t_0) e_{\pm}(t_0)$ are periodic column vectors. If there exists only one eigenvector corresponding to the repeated eigenvalue $\mu = 0$, then the general solution becomes

$$x(t) = c_+ \mathcal{P}_+(t, t_0) + c_2 [(t - t_0) \mathcal{P}_+(t, t_0) + \mathcal{P}_g(t, t_0)] \quad (47)$$

where $\mathcal{P}_g(t, t_0) = P(t, t_0) e_g(t_0)$ with $e_g(t_0)$ is a generalized eigenvector.

From above discussion, we have exponentially growing solutions iff the real part of the Floquet exponents $\Re[\mu] \neq 0$. In order to calculate Floquet exponents $\pm\mu$, it is useful to consider the eigenvalues π_{\pm} of $\mathcal{O}(t_0 + T, t_0)$ called Floquet multipliers. Since $\det \mathcal{O}(t_0 + T, t_0) = 1$, then $\pi_+ \cdot \pi_- = 1$. They are related to the Floquet exponents via $\pi_{\pm} = e^{\pm T\mu}$. In general μ and π_{\pm} are complex. Using $\pi_{\pm} = |\pi_{\pm}| e^{i\theta_{\pm}}$ and $\pi_+ \cdot \pi_- = 1$, we have

$$\mu = \frac{1}{T} [\ln |\pi_+| + i\theta_+] = -\frac{1}{T} [\ln |\pi_-| + i\theta_-].$$

Thus, $\Re[\mu] \neq 0$ iff $|\pi_{\pm}| \neq 1$.

Calculating the Floquet exponents explicitly for the problem at hand reduces to the following steps:

1. First we calculate the period T of U . The period of $U(t)$ depends on the initial amplitude of the homogeneous field $\bar{\varphi}(t_0)$ (assuming $\partial_t \bar{\varphi}(t_0) = 0$) and is given by

$$T(\bar{\varphi}_{max}) = 2 \int_{\bar{\varphi}_{min}}^{\bar{\varphi}_{max}} \frac{d\bar{\varphi}}{\sqrt{2V(\bar{\varphi}_{max}) - 2V(\bar{\varphi})}}. \quad (48)$$

In practice, we specify either $\bar{\varphi}_{max}$ or $\bar{\varphi}_{min}$. The other is found by solving $V(\bar{\varphi}_{min}) = V(\bar{\varphi}_{max})$. For $V(\varphi) = V(-\varphi)$ we have $\bar{\varphi}_{max} = \bar{\varphi}_{min}$.

2. Next we solve $\partial_t \mathcal{O}(t, t_0) = E(t) \mathcal{O}(t, t_0)$ from t_0 to $t_0 + T$ to obtain:

$$\mathcal{O}(t, t_0) = \begin{pmatrix} \delta\varphi_k^{(1)}(t_0 + T) & \delta\varphi_k^{(2)}(t_0 + T) \\ \partial_t \delta\varphi_k^{(1)}(t_0 + T) & \partial_t \delta\varphi_k^{(2)}(t_0 + T) \end{pmatrix}$$

where $\{\delta\varphi_k^{(1)}(t_0) = 1, \partial_t \delta\varphi_k^{(1)}(t_0) = 0\}$, and $\{\delta\varphi_k^{(2)}(t_0) = 0, \partial_t \delta\varphi_k^{(2)}(t_0) = 1\}$. This is equivalent to solving $\partial_t^2 \varphi_k + [k^2 + V''(\bar{\varphi})] \delta\varphi_k = 0$ for the above two sets of initial conditions from t_0 to $t_0 + T$. We have suppressed the dependence of T on $\bar{\varphi}_{max}$ to reduce clutter.

3. Last, we find the eigenvalues of $\mathcal{O}(t_0 + T, t_0)$. They are

$$\pi_{\pm}(k) = \frac{\delta\varphi_k^{(1)} + \partial_t \delta\varphi_k^{(2)}}{2} \pm \frac{\sqrt{\left\{\delta\varphi_k^{(1)} + \partial_t \delta\varphi_k^{(2)}\right\}^2 - 4}}{2} \quad (49)$$

where all quantities are evaluated at $t_0 + T$. Note that since the $\delta\varphi_k(t_0 + T)$ depends on k , the eigenvalues also depend on k . The Floquet exponents $\pm\mu$ are then given by

$$\pm\mu_k = \frac{1}{T} \ln \left[\frac{\delta\varphi_k^{(1)} + \partial_t \delta\varphi_k^{(2)}}{2} \pm \frac{\sqrt{\left\{\delta\varphi_k^{(1)} + \partial_t \delta\varphi_k^{(2)}\right\}^2 - 4}}{2} \right]. \quad (50)$$

9 Appendix B

ISW effect from nonlinear dynamics of the quintessence field

In the main body of the text, we discussed how the quintessence field fluctuations grow rapidly during the oscillatory regime. This rapid growth leads to a scale dependent growth in the gravitational potential which in turn leads to large changes in the CMB temperature anisotropies via the integrated Sachs-Wolfe (ISW) effect (see section 5.1). However, after $a_{nl} \approx 0.95$, the field perturbations become nonlinear, and the homogeneous field fragments rapidly, potentially forming long-lived, localized excitations of the field called oscillons. Our linear analysis did not include the ISW effect resulting from the nonlinear evolution of the field perturbations. In this appendix, we remedy this, by estimating the change in the CMB temperature due to:

- The change in the gravitational potential from the initial quasi/nonlinear evolution of the field perturbations.
- The time varying gravitational potential due to an isolated oscillon after it is formed.

Let us first look at the ISW effect from the initial evolution of quasi/nonlinear perturbations. A realistic calculation requires input from detailed simulations of the nonlinear field dynamics (for example, [15]). Here we concentrate on a related toy problem of calculating the ISW effect due to the collapse of a single spherical top-hat overdensity. This represents a crude approximation to the initial rapid quasi/nonlinear evolution of density perturbations associated with our model of quintessence. Later, we will evaluate the CMB temperature anisotropies from the ISW effect due to a stable oscillon like configuration. Since the universe expands very little between $a_{\text{nl}} \approx 0.95$ and $a = 1$ we will work in a Minkowski background. Our main purpose is to obtain an order of magnitude estimate and understand how the effect dependence on aspects of nonlinear evolution.

Consider a spherical mass \mathcal{M} with uniform density and a time dependent radius $R(t)$ located at \mathbf{x}_c . The gravitational potential due to this mass is given by

$$\Psi(t, \mathbf{x}) = \begin{cases} -\frac{G\mathcal{M}}{|\mathbf{x} - \mathbf{x}_c|} & |\mathbf{x} - \mathbf{x}_c| > R(t), \\ \frac{G\mathcal{M}}{2R} \left(\frac{|\mathbf{x} - \mathbf{x}_c|^2}{R^2} - 3 \right) & |\mathbf{x} - \mathbf{x}_c| \leq R(t). \end{cases}$$

The partial time derivative of this potential is

$$\partial_t \Psi(t, \mathbf{x}) = \frac{3G\mathcal{M}\dot{R}}{2R^4} u \Theta(u), \quad (51)$$

where $u = R^2(t) - |\mathbf{x} - \mathbf{x}_c|^2$ and Θ is the Heavyside function. The change in the CMB temperature due to a time varying potential (ISW term) is

$$\frac{\Delta T(\mathbf{n})}{T} = 2 \int_{t_{\text{nl}}}^{t_0} dt \partial_t \Psi(t, \mathbf{x})|_{\mathbf{x}=(t_0-t)\mathbf{n}}, \quad (52)$$

where \mathbf{n} is a unit vector that defines observer's line of sight, t_0 is the time since last scattering $\approx 0.99H_0^{-1}$ and we replace \mathbf{x} by $(t_0 - t)\mathbf{n}$ since light travels on null geodesics. We have ignored anisotropic stress here ($\Psi = \Phi$). Note that the time of integration starts at $t_{\text{nl}} \approx 0.94H_0^{-1}$ (corresponding to $a_{\text{nl}} \approx 0.95$) since nonlinear perturbations do not exist beforehand.

Let us consider “linear” collapse, ie.

$$R(t) = R_{\text{nl}} - v(t - t_{\text{nl}}), \quad (53)$$

where $0 < v < 1$. This yields

$$u = (1 - v^2)(t_+ - t)(t - t_-), \quad t_{\pm} = \frac{t_c - v(R_{\text{nl}} + vt_{\text{nl}}) \pm r_c}{1 - v^2}, \quad (54)$$

where $t_c = t_0 - \mathbf{x}_c \cdot \mathbf{n} = t_0 - |\mathbf{x}_c| \cos \theta$, $R_c = R(t_c)$, $r_c = \sqrt{R_c^2 - (1 - v^2)|\mathbf{x}_c|^2 \sin^2 \theta}$ and θ is the angle between \mathbf{n} and \mathbf{x}_c . Note that t_- and t_+ denote the time a light ray enters and

exist the spherical mass. With these definitions at hand, the integral over time in equation (52) can be evaluated analytically. In particular, if $t_{\text{nl}} < t_- < t_+ < t_0$ (and $R_c > 0$) we find

$$\frac{\Delta T(\mathbf{n})}{T} = -\frac{4G\mathcal{M}v(1-v^2)^2r_c^3}{(R_c^2 - vr_c^2)^2}. \quad (55)$$

Recalling that r_c , and R_c depend on the angle θ , the above expression shows that we should expect a cold-spot in the CMB, with an angular size $\sim R_c/|\mathbf{x}_c|$. The anisotropy is negative and maximal when $\theta = 0$, when the light ray passes through the center of the mass distribution:

$$\left.\frac{\Delta T(\mathbf{n})}{T}\right|_{\text{max}} = -\frac{4G\mathcal{M}v}{R_c}. \quad (56)$$

The result depends only on the size of the sphere R_c when the light passes through its center, the speed of collapse v and the mass \mathcal{M} . To obtain an order of magnitude estimate, we use our fiducial set of parameters. For an initial perturbation of size $R_{\text{nl}} \sim k_{\text{nl}}^{-1} \approx (0.05m)^{-1} \approx 0.02H_0^{-1}$ (see section 4.2.2 and 4.2.3), and density of order the critical density located at a distance $|\mathbf{x}_c| = 0.03H_0^{-1}$ from us (corresponding $\sim (t_0 - t_{\text{nl}})/2$), undergoing collapse with $v \sim 0.05$ (the approximate group velocity of a perturbation with wavenumber k_{nl}), we get

$$\left.\frac{\Delta T(\mathbf{n})}{T}\right|_{\text{max}} \sim -4 \times 10^{-5}, \quad (57)$$

which could provide additional constraints on our models. This signal is qualitatively similar to the Sunyaev Zel'dovich (SZ) temperature decrement from galaxy clusters [68]. However unlike \lesssim arcmin angular scale of the SZ decrement, here, the angular scale is ~ 30 degrees. As can be checked, the conditions $t_{\text{nl}} < t_- < t_+ < t_0$ and $R_c > 0$ are satisfied for the above mentioned parameters. We remind the reader that this is merely an estimate based on a spherical top-hat collapse. The actual nonlinear field evolution can be more complicated. In addition, the amplitude of the effect depends on the parameters in a non-trivial manner. Nevertheless, this exercise shows that it is indeed interesting to pursue the nonlinear evolution of the scalar field perturbations in detail.

Let us now consider the ISW effect resulting from the time varying potential associated with a single oscillon. Based on our numerical solutions (radial only), oscillons with a central field amplitude of order $M^{\ddagger\dagger}$ and width of order few m^{-1} , have a peak energy density of order m^2M^2 and an energy density profile that breathes in and out. This is in contrast with a fixed energy density profile often used in the oscillon literature. For an oscillon, crudely approximated by the a top-hat density configuration, we take

$$R_{\text{osc}}(t) \approx 2m^{-1} \left(1 + \frac{1}{4} \sin^2[mt + \phi] \right), \quad (58)$$

where for our fiducial model $m \sim 10^3 H_0$. We stress that the actual energy density profile of oscillons in our model is more complicated. The energy density profile resembles a gaussian

^{††}This M determines the scale where the scalar field potential changes shape, and should not be confused with \mathcal{M} use to denote the mass of perturbations.

when the central field amplitude is at its maximum, but becomes flatter when the field amplitude passes through zero. As a result, a calculation based on realistic oscillon profiles could be somewhat different from the estimates below.

Unlike the linear collapse case, we did not find a simple analytic expression for the ISW contribution from individual oscillons. Evaluating it numerically, we find that for our fiducial set of parameters and assuming that the energy density within oscillons $\rho \sim m^2 M^2 \sim 3H_0^2 m_{\text{pl}}^2$, we get

$$\frac{\Delta T(\mathbf{n})}{T}|_{\text{max}} \sim \pm G \mathcal{M}_{\text{osc}} m \times 10^{-2} \sim \pm 10^{-7}. \quad (59)$$

We have assumed that the oscillon is located at $|\mathbf{x}_c| \sim 0.03H_0^{-1}$. The shape of the anisotropy pattern is ring-like. The phase of the radial oscillation ϕ determines whether the center has a temperature decrement or increment. Note that the radius associated with an individual oscillon is small compared to the quasilinear linear perturbations considered earlier in this appendix (by an order of magnitude). As a result, one expects a smaller effect if one assumes that their density is still comparable to the average cosmological density. In addition, since the light crossing time is comparable to the oscillatory timescale of the energy density configuration, the ISW term undergoes cancellations, leading to a somewhat smaller ISW effect than what would be expected on purely dimensional grounds. Thus we expect the ISW contribution from individual oscillons to be smaller (and more localized) than that from the rapid collapse of nonlinear perturbations.

Before we end this appendix, we would like to comment on a caveat regarding the emergence of oscillons. Our linear analysis in the main body of the paper revealed that perturbations become non-linear at a_{nl} . While the non-linear perturbations will form oscillons eventually, it is worth asking whether we can see them as large overdensities with individual identities by today. To unambiguously understand the timescales associated with emergence of oscillons would require a full lattice simulation which is beyond the scope of this current paper. Our analysis here indicates that it is certainly worth exploring this further. While here, we have concentrated on isolated inhomogeneities, it would be interesting to look at the combined effect of a collection of such inhomogeneities (with a number density of order $(k_{\text{nl}}/2\pi)^3$) [9], which would be closer to the actual scenario.

References

- [1] A. H. Guth, Phys. Rev. D **23**, 347 (1981).
- [2] A. D. Linde, Phys. Lett. B **108**, 389 (1982).
- [3] A. Albrecht and P. J. Steinhardt, Phys. Rev. Lett. **48**, 1220 (1982).
- [4] B. Ratra and P. J. E. Peebles, Phys. Rev. D **37**, 3406 (1988).
- [5] C. Armendariz-Picon, V. F. Mukhanov, P. J. Steinhardt, Phys. Rev. Lett. **85**, 4438-4441 (2000). [astro-ph/0004134].

- [6] E. J. Copeland, M. Sami and S. Tsujikawa, *Int. J. Mod. Phys. D* **15** (2006) 1753 [arXiv:hep-th/0603057].
- [7] J. H. Traschen and R. H. Brandenberger, *Phys. Rev. D* **42**, 2491 (1990).
- [8] L. Kofman, A. D. Linde and A. A. Starobinsky, *Phys. Rev. Lett.* **73**, 3195 (1994) [arXiv:hep-th/9405187].
- [9] M. A. Amin, arXiv:1006.3075 [astro-ph.CO].
- [10] M. A. Amin, R. Easther and H. Finkel, *JCAP* **1012**, 001 (2010) [arXiv:1009.2505 [astro-ph.CO]].
- [11] R. Allahverdi, R. Brandenberger, F. Y. Cyr-Racine and A. Mazumdar, arXiv:1001.2600 [hep-th].
- [12] J. A. Frieman, C. T. Hill, A. Stebbins and I. Waga, *Phys. Rev. Lett.* **75**, 2077 (1995) [arXiv:astro-ph/9505060].
- [13] M. Khlopov, B. A. Malomed and I. B. Zeldovich, *Mon. Not. Roy. Astron. Soc.* **215** (1985) 575.
- [14] M. C. Johnson and M. Kamionkowski, *Phys. Rev. D* **78**, 063010 (2008) [arXiv:0805.1748 [astro-ph]].
- [15] M. A. Amin, R. Easther, H. Finkel, R. Flauger, M. P. Hertzberg, [arXiv:1106.3335 [astro-ph.CO]].
- [16] I. L. Bogolyubsky and V. G. Makhankov, *Pisma Zh. Eksp. Teor. Fiz.* **24**, 15 (1976).
- [17] H. Segur and M. D. Kruskal, *Phys. Rev. Lett.* **58** (1987) 747.
- [18] M. Gleiser, *Phys. Rev. D* **49**, 2978 (1994) [arXiv:hep-ph/9308279].
- [19] E. J. Copeland, M. Gleiser and H. R. Muller, *Phys. Rev. D* **52**, 1920 (1995) [arXiv:hep-ph/9503217].
- [20] J. McDonald, *Phys. Rev. D* **66**, 043525 (2002) [arXiv:hep-ph/0105235].
- [21] S. Kasuya, M. Kawasaki and F. Takahashi, *Phys. Lett. B* **559**, 99 (2003) [arXiv:hep-ph/0209358].
- [22] M. Gleiser and R. C. Howell, *Phys. Rev. Lett.* **94**, 151601 (2005) [arXiv:hep-ph/0409179].
- [23] M. Gleiser, *Int. J. Mod. Phys. D* **16**, 219 (2007) [arXiv:hep-th/0602187].
- [24] M. Hindmarsh and P. Salmi, *Phys. Rev. D* **74**, 105005 (2006) [arXiv:hep-th/0606016].
- [25] P. M. Saffin and A. Tranberg, *JHEP* **0701**, 030 (2007) [arXiv:hep-th/0610191].
- [26] E. Farhi, N. Graham, A. H. Guth, N. Iqbal, R. R. Rosales and N. Stamatopoulos, *Phys. Rev. D* **77**, 085019 (2008) [arXiv:0712.3034 [hep-th]].
- [27] G. Fodor, P. Forgacs, Z. Horvath and M. Mezei, *Phys. Lett. B* **674** (2009) 319 [arXiv:0903.0953 [hep-th]].

- [28] M. Gleiser and D. Sicilia, arXiv:0910.5922 [hep-th].
- [29] M. A. Amin, D. Shirokoff, Phys. Rev. **D81**, 085045 (2010). [arXiv:1002.3380 [astro-ph.CO]].
- [30] M. P. Hertzberg, Phys. Rev. **D82**, 045022 (2010). [arXiv:1003.3459 [hep-th]].
- [31] M. Gleiser, N. Graham and N. Stamatopoulos, Phys. Rev. D **82**, 043517 (2010) [arXiv:1004.4658 [astro-ph.CO]].
- [32] M. Gleiser, N. Graham and N. Stamatopoulos, Phys. Rev. D **83**, 096010 (2011) [arXiv:1103.1911 [hep-th]].
- [33] S. Y. Khlebnikov and I. I. Tkachev, Phys. Rev. D **56**, 653 (1997) [arXiv:hep-ph/9701423].
- [34] R. Easther and E. A. Lim, JCAP **0604**, 010 (2006) [arXiv:astro-ph/0601617].
- [35] J. F. Dufaux, A. Bergman, G. N. Felder, L. Kofman and J. P. Uzan, Phys. Rev. D **76**, 123517 (2007) [arXiv:0707.0875 [astro-ph]].
- [36] M. Y. Khlopov, Res. Astron. Astrophys. **10**, 495 (2010) [arXiv:0801.0116 [astro-ph]].
- [37] S. DeDeo, R. R. Caldwell, P. J. Steinhardt, Phys. Rev. **D67**, 103509 (2003). [astro-ph/0301284].
- [38] J. Weller, A. M. Lewis, Mon. Not. Roy. Astron. Soc. **346**, 987-993 (2003). [astro-ph/0307104].
- [39] R. Bean, O. Dore, Phys. Rev. **D69**, 083503 (2004). [astro-ph/0307100].
- [40] W. Hu, R. Scranton, Phys. Rev. **D70**, 123002 (2004). [astro-ph/0408456].
- [41] D. Rapetti, S. W. Allen, J. Weller, Mon. Not. Roy. Astron. Soc. **360**, 555-564 (2005). [astro-ph/0409574].
- [42] J. -M. Alimi, A. Fuzfa, V. Boucher, Y. Rasera, J. Courtin, P. -S. Corasaniti, Mon. Not. Roy. Astron. Soc. **401** (2010) 775. [arXiv:0903.5490 [astro-ph.CO]].
- [43] R. de Putter, D. Huterer, E. V. Linder, Phys. Rev. **D81**, 103513 (2010). [arXiv:1002.1311 [astro-ph.CO]].
- [44] R. Bean, E. E. Flanagan, M. Trodden, Phys. Rev. **D78**, 023009 (2008). [arXiv:0709.1128 [astro-ph]].
- [45] D. Bertacca, M. Bruni, O. F. Piattella and D. Pietrobon, JCAP **1102**, 018 (2011) [arXiv:1011.6669 [astro-ph.CO]].
- [46] M. Mortonson, W. Hu, D. Huterer, Phys. Rev. **D80**, 067301 (2009). [arXiv:0908.1408 [astro-ph.CO]].
- [47] E. Silverstein and A. Westphal, Phys. Rev. D **78**, 106003 (2008) [arXiv:0803.3085 [hep-th]].
- [48] L. McAllister, E. Silverstein, A. Westphal, Phys. Rev. **D82**, 046003 (2010). [arXiv:0808.0706 [hep-th]].

- [49] R. Flauger, L. McAllister, E. Pajer, A. Westphal, G. Xu, JCAP **1006**, 009 (2010). [arXiv:0907.2916 [hep-th]].
- [50] R. Kallosh and A. Linde, JCAP **1011**, 011 (2010) [arXiv:1008.3375 [hep-th]].
- [51] X. Dong, B. Horn, E. Silverstein, A. Westphal, Phys. Rev. **D84**, 026011 (2011). [arXiv:1011.4521 [hep-th]].
- [52] S. Panda, Y. Sumitomo and S. P. Trivedi, Phys. Rev. D **83**, 083506 (2011) [arXiv:1011.5877 [hep-th]].
- [53] L. J. Hall, Y. Nomura, S. J. Oliver, Phys. Rev. Lett. **95**, 141302 (2005). [astro-ph/0503706].
- [54] P. Serra, A. Cooray, D. E. Holz, A. Melchiorri, S. Pandolfi, D. Sarkar, Phys. Rev. **D80**, 121302 (2009). [arXiv:0908.3186 [astro-ph.CO]].
- [55] E. Komatsu *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **192**, 18 (2011). [arXiv:1001.4538 [astro-ph.CO]].
- [56] D. Rapetti, S. W. Allen, M. A. Amin, R. D. Blandford, Mon. Not. Roy. Astron. Soc. **375**, 1510-1520 (2007). [astro-ph/0605683].
- [57] T. M. Davis, E. Mortsell, J. Sollerman, A. C. Becker, S. Blondin, P. Challis, A. Clocchiatti, A. V. Filippenko *et al.*, Astrophys. J. **666**, 716-725 (2007). [astro-ph/0701510].
- [58] A. G. Riess, L. Macri, S. Casertano, M. Sosey, H. Lampeitl, H. C. Ferguson, A. V. Filippenko, S. W. Jha *et al.*, Astrophys. J. **699**, 539-563 (2009). [arXiv:0905.0695 [astro-ph.CO]].
- [59] R. Kessler, A. Becker, D. Cinabro, J. Vanderplas, J. A. Frieman, J. Marriner, T. MDavis, B. Dilday *et al.*, Astrophys. J. Suppl. **185**, 32-84 (2009). [arXiv:0908.4274 [astro-ph.CO]].
- [60] U. Seljak, M. Zaldarriaga, Astrophys. J. **469**, 437-444 (1996). [astro-ph/9603033].
- [61] S. Weinberg, Oxford, UK: Oxford Univ. Pr. (2008) 593 p.
- [62] W. Magnus, S. Winkler, (Interscience Publishers, New York, 1966)
- [63] L. Kofman, A. D. Linde and A. A. Starobinsky, Phys. Rev. D **56**, 3258 (1997) [arXiv:hep-ph/9704452].
- [64] G. Fodor, P. Forgacs, Z. Horvath and M. Mezei, Phys. Rev. D **79**, 065002 (2009) [arXiv:0812.1919 [hep-th]].
- [65] M. Gleiser and D. Sicilia, Phys. Rev. Lett. **101**, 011602 (2008) [arXiv:0804.0791 [hep-th]].
- [66] R. K. Sachs, A. M. Wolfe, Astrophys. J. **147**, 73-90 (1967).
- [67] C. -P. Ma, E. Bertschinger, Astrophys. J. **455**, 7-25 (1995). [astro-ph/9506072].
- [68] T. A. Marriage, V. Acquaviva, P. A. R. Ade, P. Aguirre, M. Amiri, J. W. Appel, L. F. Barrientos, E. S. Battistelli *et al.*, [arXiv:1010.1065 [astro-ph.CO]].

- [69] N. Kaiser, *Astrophys. J.* **498**, 26 (1998). [astro-ph/9610120].
- [70] W. Hu, *Astrophys. J.* **522**, L21-L24 (1999). [astro-ph/9904153].
- [71] R. Massey, J. Rhodes, A. Leauthaud, P. Capak, R. Ellis, A. Koekemoer, A. Refregier, N. Scoville *et al.*, *Astrophys. J. Suppl.* **172**, 239-253 (2007). [astro-ph/0701480].
- [72] Z. Ivezić, J. A. Tyson, R. Allsman, J. Andrew, R. Angel, f. t. L. Collaboration, [arXiv:0805.2366 [astro-ph]].
- [73] M. A. Amin, R. V. Wagoner, R. D. Blandford, *Mon. Not. Roy. Astron. Soc.* **390**, 131-142 (2008). [arXiv:0708.1793 [astro-ph]].
- [74] E. Bertschinger, P. Zukin, *Phys. Rev.* **D78**, 024015 (2008). [arXiv:0801.2431 [astro-ph]].
- [75] B. Jain and J. Khoury, *Annals Phys.* **325**, 1479 (2010) [arXiv:1004.3294 [astro-ph.CO]].
- [76] I. Laszlo, R. Bean, *Phys. Rev.* **D77**, 024048 (2008). [arXiv:0709.0307 [astro-ph]].
- [77] Y. Li and W. Hu, arXiv:1107.5120 [astro-ph.CO].
- [78] G. -B. Zhao, B. Li, K. Koyama, [arXiv:1105.0922 [astro-ph.CO]].
- [79] B. Li, J. D. Barrow, [arXiv:1010.3748 [astro-ph.CO]].
- [80] B. Li, D. F. Mota, J. D. Barrow, *Astrophys. J.* **728**, 109 (2011). [arXiv:1009.1400 [astro-ph.CO]].
- [81] A. Arvanitaki, S. Dimopoulos, S. Dubovsky, N. Kaloper and J. March-Russell, *Phys. Rev. D* **81**, 123530 (2010) [arXiv:0905.4720 [hep-th]].
- [82] D. J. E. Marsh, P. G. Ferreira, *Phys. Rev.* **D82**, 103528 (2010). [arXiv:1009.3501 [hep-ph]].